




SINGLE APPLICATION FLEXIBILITY FOR SEASON LONG CONTROL – SOYBEAN

DUAL MODE OF ACTION PRODUCT PROVIDING BROAD-SPECTRUM ACTIVITY ON KEY SOYBEAN PATHOGENS

Frogeye Leaf Spot Trial - 2014

Mississippi State University
Delta Research and Extension Center – Stoneville, MS




All fungicide treatments applied July 17th (R4) with non-ionic surfactant at 0.25% v/v.



White Mold Trial - 2015

University of Wisconsin
Hancock Agricultural Experiment Station – Hancock, WI



Key Benefits

- 2 modes of action, flutriafol (Triazole) + fluoxastrobin (Strobilurin) for resistance management especially on tough diseases such as frogeye leaf spot.
- Highly systemic in xylem.
- Systemicity + long residual providing application flexibility.
- Plant health effects including: decreased ethylene production, increased carbon assimilation, increased water and nutrient uptake efficiency.
- Improved stress tolerance, greener healthier plants allowing maximum yield.

Application Guidelines

- R1-R5 – 5 oz/a
- Adjuvant use: Including NIS 0.25% v/v or an oil based adjuvant 0.5-1% v/v may improve spray coverage and canopy penetration, if tank mixed follow adjuvant requirements for tank mix partner.

Application Recommendations

Control: Alternaria leaf spot, Anthracnose, Brown spot, Cercospora blight, and Purple seed stain, Frogeye leaf spot, Pod and Stem blight, Powdery mildew, Rhizoctonia aerial blight, Rust.

4-6 oz/A

Suppression: Sclerotinia stem rot, White mold.

12 Hour REI, 30 Day PHI (seed harvest).

Apply using nozzles delivering thorough coverage to provide adequate disease control. For ground application equipment apply in a minimum of 10 gal/a. For aerial application, use aircraft spray equipment in a minimum of 2 gal/a. Can be tank mixed with commonly used herbicides, fungicides, insecticides and foliar nutrients. However, compatibility should be determined by jar testing.

Special Information

Do not apply more than 12 oz/a per acre per year. Do not apply more than 0.132 lb ai/a fluoxastrobin and 0.227 lb ai/a flutriafol per year. Apply no later than R5 (beginning seed). Do not make more than two applications per year. The minimum retreatment interval is 14 days. Do not feed forage or hay to animals or permit animals to graze.

Rotational Crop Guideline

Soybeans, Corn, Wheat, Peanuts	Anytime
Sugar Beets	30 days
Sweet Corn, Cotton	180 days

Always read and follow label directions. FMC and Preemptor are trademarks of FMC Corporation or an affiliate. Approach and Approach Prima are trademarks of E.I. duPont De Nemours and Co. Headline and Priaxor are trademarks of BASF SE. Quadris, Quadris TOP and Quilt Xcel are trademarks of Syngenta Group Company. Stratego YLD and Proline are trademarks of Bayer CropScience LP ©2015 FMC Corporation. All rights reserved.